

HERPC Hull & East Riding Prescribing Committee **NEWSLETTER NOVEMBER 2020**

For further information about HERPC documents go to:

<http://www.hey.nhs.uk/herpc>

For Joint Formulary go to

<http://www.hey.nhs.uk/herpc/joint-formulary.htm>

Traffic Light Classification within HERPC:

RED – specialist prescriber only

AMBER – prescribed in accordance with approved shared care framework

BLUE (Guideline or Specialist Led) – prescribed on advice of specialist or in line with national / local guideline

GREEN – other items listed on formulary suitable for initiation and prescribing by any prescriber

ADDITIONS TO RED LIST

Tropicamide/Phenylephrine/Lidocaine (Mydrane®) for Mydriasis & Intraocular anaesthesia

Trametinib Low Grade Ovarian Cancer

Atezolizumab TA 638 Atezolizumab with carboplatin and etoposide for untreated extensive-stage small-cell lung cancer

Atezolizumab with nab-paclitaxel TA 639 Atezolizumab with nab-paclitaxel for untreated PD-L1-positive, locally advanced or metastatic, triple-negative breast cancer

Lorlatinib TA628 Lorlatinib for previously treated ALK-positive advanced non-small-cell lung cancer

Gilteritinib TA 642 Gilteritinib for treating relapsed or refractory acute myeloid leukaemia

Avelumab with axitinib TA645 Avelumab with axitinib for untreated advanced renal cell carcinoma

Gemtuzumab ozogamicin TA545 Untreated acute myeloid leukaemia

Encorafenib and Cetuximab Advanced Disease with B-Raf V600E mutation

Darolutamide Castrate resistant high risk non metastatic prostate cancer with LHRH antagonist injections

Avelumab Monotherapy for first line maintenance treatment of adult patients with locally advanced or metastatic urothelial carcinoma

NEW & UPDATED SCF

SCF Sandostatin analogues for management of high output stomas (off label use of licensed medicines)

Riluzole SCF

NEW & UPDATED GUIDELINES/SPECIALIST LED MEDICINES

Tinzaparin for use in paediatrics at HUTH

Lenzetto Estradiol Spray Hormone Replacement Therapy (HRT) for oestrogen deficiency symptoms in postmenopausal women
(in women at least 6 months since last menses or surgical menopause, with or without a uterus).

Lurasidone Schizophrenia in adults and adolescents over 13yrs of age

Thickener Prescribing Guide & Thickeners and Dysphagia in Carehomes

BPSD Pathway – replace with NICE Dementia Guidance

High Output Stoma Guideline

GREEN DRUGS

None this month